

Universität Bonn

Physikalisches Institut

New $N=1$ Extended Superconformal Algebras with Two and Three Generators

(to be published in Int. Jour. of Mod. Phys. **A**)

R. Blumenhagen, W. Eholzer, A. Honecker, R. Hübel

Abstract

In this paper we consider extensions of the super Virasoro algebra by one and two super primary fields. Using a non-explicitly covariant approach we compute all $\mathcal{SW}$ -algebras with one generator of dimension up to 7 in addition to the super Virasoro field. In complete analogy to $\mathcal{W}$ -algebras with two generators most results can be classified using the representation theory of the super Virasoro algebra. Furthermore, we find that the $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ -algebra can be realized as a subalgebra of $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$ at $c = \frac{10}{7}$. We also construct some new $\mathcal{SW}$ -algebras with three generators, namely $\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, \frac{5}{2})$, $\mathcal{SW}(\frac{3}{2}, 2, 2)$ and $\mathcal{SW}(\frac{3}{2}, 2, \frac{5}{2})$.

Post address:
Nußallee 12
W-5300 Bonn 1
Germany
e-mail: unp039@ibm.rhrz.uni-bonn.de

BONN-HE-92-02
hep-th/9207072
Bonn University
January 1992
ISSN-0172-8733

1. Introduction

Extended conformal and superconformal algebras have gained intensive interest in theoretical physics because they can be considered as symmetry algebras of conformal quantum field theories (CFT). One hopes to describe all rational conformal quantum field theories (RCFT) as minimal models of extended chiral algebras [1], so-called $\mathcal{W}$ -algebras.

The first $\mathcal{W}$ -algebras were constructed by Zamolodchikov in 1985 [2]. He extended the Virasoro algebra by primary fields up to dimension three. Two of these $\mathcal{W}$ -algebras have the typical property that normal ordered products appear. We denote the extension of the Virasoro algebra by primary fields of conformal dimensions $\delta_1, \dots, \delta_n$ by $\mathcal{W}(2, \delta_1, \dots, \delta_n)$. In the last years, several methods have been developed to find new $\mathcal{W}$ -algebras. One approach uses the free field construction based on Lie algebras and their infinite dimensional extensions, the Kac-Moody algebras [3][4][5].

In this paper we use a different method, constructing the algebra explicitly. Using this method a whole bunch of $\mathcal{W}$ -algebras has been constructed [6][7][8][9][10][11]. In [10][11] a constructive algorithm has been presented which is based on a structural theorem about chiral $SU(1, 1)$ invariant algebras [12]. It has been shown in [9] that $\mathcal{W}$ -algebras with two generators can only exist for generic values of the central charge, if the conformal dimension of the additional primary field is contained in $\{\frac{1}{2}, 1, \frac{3}{2}, 2, 3, 4^{[7][6]}, 6^{[8]}\}$. In [10][11] it has been shown that several other $\mathcal{W}(2, \delta)$ -algebras exist for finitely many values of the central charge c . The possible c values are essentially determined by the representation theory of the Virasoro algebra.

RCFT's describe two dimensional statistical models at second order phase transitions. Furthermore there are statistical models which possess superconformal invariance [13] at criticality. These models are described by minimal models of the super Virasoro algebra. Similar to the Virasoro algebra, the minimal series is bounded from above by $c = \frac{3}{2}$. In order to get minimal models with higher values of the central charge one investigates extensions of the super Virasoro algebra, called super $\mathcal{W}$ -algebras.

In this paper we consider extensions of the super Virasoro algebra by one and two super primary fields. In general we denote the extension of the super Virasoro algebra by super primary fields of dimension $d_1, \dots, d_n$ by $\mathcal{SW}(\frac{3}{2}, d_1, \dots, d_n)$. Some $\mathcal{SW}(\frac{3}{2}, d)$ -algebras have already been calculated in [14][15][16][17]. As far as one knows $\mathcal{SW}(\frac{3}{2}, d)$ -algebras exist generically only for $d \in \{\frac{1}{2}, 1, \frac{3}{2}, 2\}$. For $d \neq 2$ these algebras are super Lie algebras. In [17] all $\mathcal{SW}(\frac{3}{2}, d)$ -algebras up to $d = \frac{7}{2}$ have been constructed using the super conformal bootstrap method. Here we extend this series of algebras up to $d = 7$ using the same methods as presented in [10]. Superconformal covariance is implemented by considering Jacobi identities involving the generator $G(z)$ of pure super transformations. This algorithm differs from the explicitly covariant one presented in [18]. Nevertheless these two algorithms seem to be equivalent. For $d \geq \frac{5}{2}$ the $\mathcal{SW}(\frac{3}{2}, d)$ -algebras exist only for discrete values of c . Most of these values can be classified in complete analogy to the $\mathcal{W}(2, \delta)$ -algebras, essentially using representation theory of the super Virasoro algebra. Furthermore, we find that the $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ -algebra can be realized as a subalgebra of the $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$ -algebra for $c = \frac{10}{7}$. A similar construction has been carried out in [19] where it has been shown that $\mathcal{SW}(\frac{3}{2}, \frac{7}{2})$ is contained in $\mathcal{SW}(\frac{3}{2}, \frac{3}{2})$ for $c = \frac{7}{5}$.

This paper is organized as follows: In the next chapter we review a general theorem about chiral $SU(1,1)$ invariant algebras. In the third chapter we give the general outline for the construction of $\mathcal{SW}$ -algebras in the non-explicitly covariant approach. Chapter four contains our results on $\mathcal{SW}(\frac{3}{2}, d)$ -algebras. Afterwards we discuss these results in chapter five. We continue with our results concerning $\mathcal{SW}(\frac{3}{2}, d_1, d_2)$ -algebras in chapter six. In chapter seven we draw some conclusions from the results obtained.

2. General theorems about $SU(1,1)$ invariant algebras

This chapter is devoted to a short review of well known results concerning algebras of local chiral fields. We will only give a brief summary; for proofs as well as for details we refer to [20][10].

Let $\mathcal{F}$ be the algebra of local chiral fields of a conformal field theory defined on 2-dimensional spacetime (with compactified space). Because of $SU(1,1)$ -invariance $\mathcal{F}$ carries a natural grading by the conformal dimension and is spanned by the non-derivative (i. e. quasiprimary) fields together with their derivatives. In $\mathcal{F}$ the operation of building normal ordered products (NOPs) is defined (see below).

We define the Fourier decomposition of a left chiral field by $\phi(z) = \sum_{n-d(\phi) \in \mathbb{Z}} z^{n-d(\phi)} \phi_n$. We call the Fourier-components ϕ_n the ‘modes’ of ϕ .

Denote the vacuum of the theory by $|v\rangle$. Requiring the regularity of $\phi(z)|v\rangle$ at the origin implies

$$\phi_n |v\rangle = 0 \quad \forall n < d(\phi) \quad (2.0)$$

It is well known that the modes of the energy momentum tensor satisfy the Virasoro algebra

$$[L_m, L_n] = (n - m)L_{m+n} + \frac{c}{12}(n^3 - n)\delta_{n+m,0} \quad (2.1)$$

with central charge c . A primary field ϕ of conformal dimension d is characterized by the commutator of its modes with the Virasoro algebra (2.1):

$$[L_m, \phi_n] = (n - (d - 1)m)\phi_{n+m} \quad (2.2)$$

Iff a field ϕ satisfies (2.2) for $m \in \{-1, 0, 1\}$ this field is called ‘quasiprimary’.

As was shown by W. Nahm [20][12] locality and invariance under rational conformal transformations already impose severe restrictions on the commutator of two quasiprimary chiral fields:

Let $\{\phi_i \mid i \in I\}$ be a set of non-derivative fields of integer or half integer dimensions $d(\phi_i) = d_i$, which together with their derivatives span $\mathcal{F}$. Then the mode algebra of the Fourier components of left chiral fields has the form

$$[\phi_{i,m}, \phi_{j,n}]_{\pm} = \sum_{k \in I} C_{\phi_i \phi_j}^{\phi_k} p_{ijk}(m, n) \phi_{k, m+n} + d_{ij} \delta_{n, -m} \binom{n + d_i - 1}{2d_i - 1} \quad (2.3)$$

Here the d_{ij} describe the normalization of the two point functions and the $C_{\phi_i\phi_j}^{\phi_k}$ are the coupling constants between three quasiprimary fields. The p_{ijk} are universal polynomials depending only on the dimensions of the fields ϕ_i, ϕ_j and ϕ_k ; their explicit form is presented in [10].

In addition to their Lie bracket structure the algebra $\mathcal{F}$ admits another important operation, namely forming normal ordered products (NOPs) of chiral fields. Usually the NOP of two chiral fields ϕ, χ is defined in terms of Fourier components as follows

$$N(\phi, \chi)_n := \epsilon_{\phi\chi} \sum_{k < d(\chi)} \phi_{n-k} \chi_k + \sum_{k \geq d(\chi)} \chi_k \phi_{n-k} \quad (2.4)$$

$\epsilon_{\phi\chi}$ is defined as -1 if ϕ and χ are both fermions and 1 otherwise.

In this form it occurs in the regular part of the OPE of ϕ and χ , but it is not a non-derivative field, so that e.g. equation (2.3) cannot be used to gain any information about its commutator with other fields. For the NOP to be ‘well behaved’ under $SU(1,1)$ -transformations we have to add some counterterms to $N(\phi, \chi)$ defined in (2.4):

With the assumptions and notations from above we define the normal ordered product of two chiral fields by [12]

$$\mathcal{N}(\phi_j, \partial^n \phi_i) := N(\phi_j, \partial^n \phi_i) - \left(\sum_{r=1}^n \alpha_{ij}^r \partial^r N(\phi_j, \partial^{n-r} \phi_i) + \sum_k \beta_{ij}^k(n) C_{\phi_i\phi_j}^{\phi_k} \partial^{\gamma_{ij}^k(n)} \phi_k \right) \quad (2.5)$$

Where $\alpha_{ij}^r, \beta_{ij}^k(n), \gamma_{ij}^k(n)$ are polynomials depending only on the conformal dimensions of the involved fields (for more details see [10]). This definition yields a quasiprimary field of conformal dimension $d_i + d_j + n$.

Since the field content of $\mathcal{F}$ is infinite one introduces the notion of non-composite, ‘simple’ fields. To be more precise, if a basis for $\mathcal{F}$ can be obtained from a set of fields $\{\phi_i\}_{i \in J}$ using the operations given above (forming normal ordered products and derivatives) we will say that the fields $\{\phi_i\}_{i \in J}$ generate $\mathcal{F}$. If all fields in $\{\phi_i\}_{i \in J}$ are quasiprimary and orthogonal to all normal ordered products we will call these fields ‘simple’. A set of simple fields will be called a simple set.

The commutators of normal ordered products are completely determined by the commutators of the simple fields involved. This means that the whole Lie algebra structure of the mode algebra of $\mathcal{F}$ is already fixed by the commutation relations of the simple fields. The coupling constants of the simple fields determine all other coupling constants.

In general, $\mathcal{W}$ -algebras are not Lie algebras since the commutators do not close linearly in the fields. If the OPE corresponding to the mode algebra is required to be associative, the mode algebra has to fulfil all possible Jacobi identities. In order to ensure the validity of all Jacobi identities it is sufficient to verify that only those involving simple fields (besides L) are satisfied. Looking at the corresponding OPE one notices that it is sufficient to

show that the factors in front of primary fields vanish. This leads to equations among the coupling constants connecting three simple fields.

3. General definitions and theorems about $\mathcal{SW}$ -algebras

In this chapter we adapt the results of the conformal case to the superconformal one. To this end, we recall the definitions of super fields, super primaries and super Jacobi identities. Furthermore, we set up some conventions and establish some simple relations between various structure constants arising from superconformal covariance.

The super Virasoro algebra is the extension of the Virasoro algebra by a primary field G of dimension $\frac{3}{2}$. Using the normalization $d_{GG} = \frac{2c}{3}$, the commutation relations take the following form:

$$\begin{aligned} [L_m, L_n] &= (n-m)L_{m+n} + \frac{c}{12}(n^3-n)\delta_{n+m,0} \\ [L_m, G_n] &= (n-\frac{1}{2}m)G_{m+n} \\ [G_m, G_n]_+ &= 2L_{m+n} + \frac{c}{3}(m^2-\frac{1}{4})\delta_{m+n,0}, \end{aligned} \tag{3.1}$$

which can also be obtained from the theorem of chapter 2. This algebra is known as the central extension of the algebra formed by the generators of superconformal transformations in super space Z consisting of the points (z, θ) , where θ is a Grassmannian variable. The two fields can be composed to the super Virasoro field $\mathcal{L} = \frac{1}{2}G + \theta L$, defined on Z . A field $\Phi = \phi + \theta\psi$ is called super primary of dimension $d = d(\Phi)$, iff ϕ and ψ are Virasoro primaries of conformal dimension d and $d + \frac{1}{2}$ respectively and

$$\begin{aligned} [G_n, \phi_m]_{\pm} &= \mathcal{C}_{G\phi}^{\psi} p_{\frac{3}{2}, d, d+\frac{1}{2}}(n, m) \psi_{n+m} = \mathcal{C}_{G\phi}^{\psi} \psi_{n+m} \\ [G_n, \psi_k]_{\pm} &= \mathcal{C}_{G\psi}^{\phi} p_{\frac{3}{2}, d+\frac{1}{2}, d}(n, k) \phi_{n+k} = \frac{\mathcal{C}_{G\psi}^{\phi}}{2d} (k - (2d-1)n) \phi_{n+k}, \end{aligned} \tag{3.2}$$

with ϕ and ψ super partners of each other; here $\mathcal{C}_{G\phi}^{\psi}$ and $\mathcal{C}_{G\psi}^{\phi}$ are certain structure constants and we have inserted the universal polynomials p_{ijk} . In perfect analogy to the quasiprimary case a field $\phi + \theta\psi$ is called super quasiprimary iff ϕ and ψ are Virasoro quasiprimaries and (3.2) holds for $n \in \{\frac{1}{2}, -\frac{1}{2}\}$. This definition is equivalent to the covariant definition of super primary fields [21].

Let $\mathcal{A} = \{G, L, \phi_1, \psi_1, \dots, \phi_n, \psi_n\}$ be a simple set with super primaries $\Phi_i = \phi_i + \theta\psi_i$. We will call the algebra generated by $\mathcal{A}$ a $\mathcal{SW}(\frac{3}{2}, d(\Phi_1), \dots, d(\Phi_n))$ and the fields $\Phi_1, \dots, \Phi_n$ additional fields.

To ensure the existence of $\mathcal{SW}(\frac{3}{2}, d_1, \dots, d_n)$ super Jacobi identities have to be checked which are of the general form:

$$(\epsilon_{\chi_i \chi_k} [[\chi_{i,p}, \chi_{j,q}]_{\pm}, \chi_{k,r}]_{\pm})_{cycl.} = 0, \tag{3.3}$$

where χ_i, χ_j and χ_k are components (ϕ or ψ) of super fields. In the sequel we will denote the factor in front of the field $\chi_{l,p+q+r}$ on the left hand side of (3.3) by $(\chi_{i,p}, \chi_{j,q}, \chi_{k,r}, \chi_{l,p+q+r})$. The simplest example for $\mathcal{SW}$ -algebras is the super Virasoro algebra (3.1) since it obeys the super Jacobi identities (3.3), which is also a super Lie algebra.

The identities (3.3) will lead to relations involving different structure constants. Some of these identities hold in general, already determined by the supersymmetric structure. This illustrates the fact that we are working with a non-explicitly covariant approach, dealing with components of super fields rather than with super fields themselves.

These relations between various structure constants can be derived from simple Jacobi identities involving the component G of the super Virasoro field. In order to give some examples we fix the normalization of the fields first:

$$\begin{aligned} d_{\phi_i \phi_i} &= \frac{(-1)^{2d(\phi_i)+1}}{2d(\phi_i) + 1} \frac{c}{d(\phi_i)} \\ d_{\psi_i \psi_i} &= \frac{c}{d(\psi_i)} \end{aligned} \quad (3.4)$$

Note that in the covariant approach this leads to the standard normalization of super fields [18].

Then we can e.g. determine the structure constants $C_{G\phi}^\psi$ and $C_{G\psi}^\phi$ in (3.2):

$$C_{G\phi_i}^{\psi_i} = \frac{1}{2d_i} C_{G\psi_i}^{\phi_i} \quad \text{and} \quad (C_{G\phi_i}^{\psi_i})^2 = 1 \quad (3.5)$$

The first of these relations between the different structure constants is a direct consequence of the normalization of the fields ϕ_i and ψ_i and the fact that they are simple fields. The other relation is easily seen for $d_i \in \mathbb{N}$ by the following argument (the case $d_i \in \mathbb{N} + \frac{1}{2}$ is treated similarly): Consider the following Jacobi identity:

$$([[G_r, G_s]_{\pm}, \phi_{i,m}]_{\pm})_{cycl.} = 0$$

Because $(G_r, G_s, \phi_{i,m}, \phi_{i,r+s+m})$ has to be zero we obtain:

$$\begin{aligned} &C_{GG}^L C_{L\phi_i}^{\phi_i} p_{2,d_i,d_i}(r+s, m) - \\ &C_{G\phi_i}^{\psi_i} C_{\psi_i G}^{\phi_i} (p_{\frac{3}{2},d_i,d_i+\frac{1}{2}}(s, m) p_{d_i+\frac{1}{2},\frac{3}{2},d_i}(s+m, r) + p_{d_i,\frac{3}{2},d_i+\frac{1}{2}}(m, r) p_{d_i+\frac{1}{2},\frac{3}{2},d_i}(m+r, s)) = 0 \end{aligned}$$

using $C_{\psi_i G}^{\phi_i} = 2d_i C_{G\phi_i}^{\psi_i}$, $C_{GG}^L = 2$ and $C_{L\phi_i}^{\phi_i} = d_i$, one obtains:

$$(C_{G\phi_i}^{\psi_i})^2 = 1$$

In this chapter we will choose the positive root, so that $C_{G\phi_i}^{\psi_i} = 1$.

Furthermore, all possible coupling constants between three super primary fields $(\phi_i, \psi_i) \times (\phi_j, \psi_j) \rightarrow (\phi_k, \psi_k)$ are determined by a single coupling constant of the components:

i) $d_i + d_j + d_k \in \mathbb{N} + \frac{1}{2}$:

$$C_{\psi_i \phi_j}^{\phi_k} = 2d_k C_{ij}^k, \quad C_{\phi_i \psi_j}^{\phi_k} = (-1)^{2d_i+1} 2d_k C_{ij}^k, \quad C_{\psi_i \psi_j}^{\psi_k} = (-1)^{2d_i+1} (\sigma_{ijk} + \frac{1}{2}) C_{ij}^k \quad (3.6a)$$

with $C_{ij}^k = C_{\phi_i \phi_j}^{\psi_k}$

ii) $d_i + d_j + d_k \in \mathbb{N}$:

$$C_{\psi_i \phi_j}^{\psi_k} = \frac{h_{ikj}}{2d_k} \hat{C}_{ij}^k, \quad C_{\psi_i \psi_j}^{\phi_k} = (-1)^{2d_i+1} h_{ijk} \hat{C}_{ij}^k, \quad C_{\phi_i \psi_j}^{\phi_k} = (-1)^{2d_i} \frac{h_{jki}}{2d_k} \hat{C}_{ij}^k \quad (3.6b)$$

with $\hat{C}_{ij}^k = C_{\phi_i \phi_j}^{\phi_k}$

and $\sigma_{ijk} = d_i + d_j + d_k - 1$, $h_{ijk} = d_i + d_j - d_k$

For a proof consider the condition

$$(G_l, \phi_{i,m}, \phi_{j,n}, \psi_{k,l+m+n}) = 0.$$

This yields an equation in the coupling constants

$$C_{\psi_i \phi_j}^{\psi_k}, \quad C_{\phi_i \psi_j}^{\psi_k}, \quad C_{\phi_i \phi_j}^{\phi_k}.$$

Setting $l = \frac{1}{2}$, $m = d_j - d_k$, $n = 1 - d_j$ and $l = -\frac{1}{2}$, $m = 1 - d_i$, $n = d_i - d_k$ one obtains two equations. Combining these equations proves the first two relations. The other relations are obtained by studying

$$\begin{aligned} (G_l, \phi_{j,m}, \psi_{i,n}, \phi_{k,l+m+n}) &= 0 \\ (G_l, \psi_{i,m}, \psi_{j,n}, \phi_{k,l+m+n}) &= 0 \\ (G_l, \psi_{j,m}, \phi_{i,n}, \psi_{k,l+m+n}) &= 0. \end{aligned}$$

Remark 3.1: If one assumes that a basis of super quasiprimary fields exists (e.g. in the covariant approach) the formulae given above hold also for super quasiprimary fields since only $l \in \{\frac{1}{2}, -\frac{1}{2}\}$ is needed for the proof. The same formulae can be deduced in the explicitly covariant approach [18].

Remark 3.2: Because of $C_{ij}^k = (-1)^{[d_i]+[d_j]+[d_k+\frac{1}{2}]} C_{ji}^k$ and $\hat{C}_{ij}^k = (-1)^{[d_i]+[d_j]+[d_k]} \hat{C}_{ji}^k$ one obtains for $(\phi_i, \psi_i) = (\phi_j, \psi_j)$:

- i) $\hat{C}_{ii}^k \neq 0 \Rightarrow d_k \in 2\mathbb{N}$
- ii) $C_{ii}^k \neq 0 \Rightarrow d_k + \frac{1}{2} \in 2\mathbb{N}$

In the explicitly covariant formulation [18] it is evident that in the singular part of the OPE of a super primary field Φ_i with Φ_j no normal ordered products of these super

fields appear. In the non-explicitly covariant approach this may happen only if the field $\mathcal{N}(\phi_i, \phi_j)$ appears in the commutator $[\psi_i, \psi_j]_{\pm}$. In this case either $\mathcal{N}(\phi_i, \phi_j)$ must have a vanishing primary projection and can be replaced by a linear combination of the other fields of dimension $d_i + d_j$ or the relevant coupling constant has to vanish. Denote the primary projection of ϕ by $\mathcal{P}\phi$. Then one has

$$\mathcal{PN}(\phi_i, \phi_j) \neq 0 \quad \Rightarrow \quad \mathcal{C}_{\psi_i \psi_j}^{\mathcal{N}(\phi_i, \phi_j)} = 0 \quad (3.7)$$

Since

$$[\psi_{i,m}, \psi_{j,n}]_{\pm} = \mathcal{C}_{\psi_i \psi_j}^{\mathcal{N}(\phi_i, \phi_j)} \mathcal{N}(\phi_i, \phi_j)_{m+n} + \dots ,$$

applying $G_{\frac{1}{2}}$ to $[\psi_{i,m}, \psi_{j,n}]_{\pm}$ yields

$$\begin{aligned} & \mathcal{C}_{G\mathcal{N}(\phi_i, \phi_j)}^{\mathcal{N}(\phi_i, \psi_j)} \mathcal{C}_{\psi_i \psi_j}^{\mathcal{N}(\phi_i, \phi_j)} \mathcal{N}(\phi_i, \psi_j)_{n+m+\frac{1}{2}} + \mathcal{C}_{G\mathcal{N}(\phi_i, \phi_j)}^{\mathcal{N}(\psi_i, \phi_j)} \mathcal{C}_{\psi_i \psi_j}^{\mathcal{N}(\phi_i, \phi_j)} \mathcal{N}(\psi_i, \phi_j)_{n+m+\frac{1}{2}} + \dots \\ &= [G_{\frac{1}{2}}, [\psi_{i,m}, \psi_{j,n}]_{\pm}]_{\pm} \\ &= (m - d_i + \tfrac{1}{2})[\phi_{i,m+\frac{1}{2}}, \psi_{j,n}]_{\pm} + (n - d_j + \tfrac{1}{2})[\psi_{i,n+\frac{1}{2}}, \phi_{j,m}]_{\pm}, \end{aligned}$$

where the dots stand for fields which are linearly independent from $\mathcal{N}(\phi_i, \psi_j)$ and $\mathcal{N}(\psi_i, \phi_j)$. Because $\mathcal{C}_{G\mathcal{N}(\phi_i, \phi_j)}^{\mathcal{N}(\phi_i, \psi_j)}$ and $\mathcal{C}_{G\mathcal{N}(\phi_i, \phi_j)}^{\mathcal{N}(\psi_i, \phi_j)}$ are nonzero and in the commutators in the last line the field $\mathcal{N}(\phi_i, \psi_i)$ does not occur $\mathcal{C}_{\psi_i \psi_j}^{\mathcal{N}(\phi_i, \phi_j)}$ has to vanish.

Remark 3.3: We use the following normalization in our calculations instead of (3.4):

$$d_{\phi_i \phi_i} = \frac{c}{d(\phi_i)} \quad \text{and} \quad d_{\psi_i \psi_i} = \frac{c}{d(\psi_i)} \quad (3.4')$$

While the normalization (3.4) is more convenient for proofs this one is more convenient for practical calculations (in this normalization the d -matrices almost factorize completely into linear factors (cf. appendix A and C)).

Now equations (3.5) and (3.6) have to be modified slightly. One obtains e.g.:

$$C_{G\phi_i}^{\psi_i} = \frac{2d_i + 1}{2d_i} (-1)^{2d_i+1} C_{G\psi_i}^{\phi_i} \quad \text{and} \quad (C_{G\phi_i}^{\psi_i})^2 = (-1)^{2d_i+1} (2d_i + 1) \quad (3.5')$$

$$(C_{\psi_i \phi_j}^{\phi_k})^2 = (2d_k)^2 \frac{(-1)^{2d_i+2d_k} (2d_k + 1)}{2d_i + 1} (C_{ij}^k)^2 \quad (3.6')$$

After this general setup we present an algorithm for the explicit constuction of $\mathcal{SW}$ -algebras. Let $\mathcal{A} = \{G, L, \phi_1, \psi_1, \dots, \phi_n, \psi_n\}$ be a simple set and $\Phi_i = \phi_i + \theta\psi_i$ super primaries of dimension d_i .

Then one proceeds as follows:

First write down all linearly independent NOP's which may appear in the commutator of the simple additional fields. The algorithm used is based on the following facts. Let ϕ be

a quasiprimary NOP of dimension δ . Then the quasiprimary projection of $\partial\phi = [L_1, \phi]$ is zero and leads to a linear equation in quasiprimary NOPs of dimension $\delta + 1$. Such equations can be used to reduce the set of all possible NOPs of dimension $\delta + 1$ to a basis. Formally, this can be described by assigning to each quasiprimary field of dimension δ a coloured partition $p(\delta)$. The linear equation obtained by the quasiprimary projection can be written in terms of partitions of $\delta + 1$. Thus the main step of the algorithm is to delete for any $p(\delta)$ an arbitrary $p(\delta + 1)$ that can be obtained by adding 1 to some element of $p(\delta)$. For details of the algorithm as well as for a proof and examples we refer the reader to [22].

Secondly, calculate all structure constants appearing in these commutators. The structure constants connecting three additional super primary fields remain as free parameters.

Finally, one has to check all Jacobi identities with additional super primary fields only. This will in general lead to conditions for the free coupling constants and the central charge.

Note that the coupling constants containing one G are determined by (3.5) and the coupling constants between the superconformal families are determined by a single coupling constant (cf. (3.6)).

In order to check the validity of the Jacobi identities it is sufficient to ensure that all factors in front of the primary fields vanish. Primary fields include the additional simple fields and perhaps fields built from NOPs of two additional fields and G, L . If the coupling constants of the additional simple fields vanish, such NOPs cannot turn up in the Jacobi identities and one has to check the factors in front of the additional simple fields only. If some coupling constants are nonzero one has to calculate which primary projections of NOPs of additional simple fields do not vanish and to ensure that the coefficients in front of these fields are zero. This is primarily a practical problem since the calculation of these factors is very complicated. Therefore we checked the factors in front of the simple fields only. In fact, there is no example showing that these conditions are not sufficient (in all previously known cases we obtained the same results). Indeed there are some examples where non-simple primary fields occur in the Jacobi identities and do not lead to further restrictions (see for example $\mathcal{SW}(\frac{3}{2}, 2)$ in the next chapter).

For the special case of $\mathcal{SW}$ -algebras with two and three generators this general outline reduces to the following.

While for $\mathcal{SW}$ -algebras with two generators there remain two free parameters (c and the self-coupling) there are five free parameters (c and four coupling constants) for $\mathcal{SW}$ -algebras with three generators. The coupling constants C_{ii}^j vanish if $d_j \in 2\mathbb{N} + 1$ or $d_j \in 2\mathbb{N} + \frac{1}{2}$ (cf. remark 3.2). In almost all calculations we verified that the Jacobi identities with $\bar{L}$ or G involved are satisfied automatically, if one uses (3.5) and (3.6). In one

superconformal family four Jacobi identities remain to be checked:

$$\begin{aligned}
[[\phi_m, \phi_n]_{\pm}, \phi_k]_{\pm, cycl.} &= 0 \\
[[\phi_m, \phi_n]_{\pm}, \psi_k]_{\pm, cycl.} &= 0 \\
[[\psi_m, \psi_n]_{\pm}, \phi_k]_{\pm, cycl.} &= 0 \\
[[\psi_m, \psi_n]_{\pm}, \psi_k]_{\pm, cycl.} &= 0
\end{aligned}$$

For $\mathcal{SW}(\frac{3}{2}, d_1, d_2)$ -algebras there are six Jacobi identities involving two fields of one additional superconformal family and one of the second family, such that altogether 20 Jacobi identities have to be verified.

We denote the coupling of Φ_i with Φ_j to Φ_k by $C_{\Phi_i \Phi_j}^{\Phi_k}$, where

$$C_{\Phi_i \Phi_j}^{\Phi_k} = \begin{cases} C_{\phi_i \phi_j}^{\phi_k} & , d(\Phi_k) \in \mathbb{N} \\ C_{\phi_i \phi_j}^{\psi_k} & , d(\Phi_k) \in \mathbb{N} + \frac{1}{2} \end{cases} \quad (3.6)$$

If there exist null fields for a possible discrete value of the central charge one has to ensure that either all further calculations are performed without these null fields and fixed central charge or that they are performed generically.

4. Explicit results on $\mathcal{SW}(\frac{3}{2}, d)$ -algebras

In this chapter we present our results on explicit constructions of several $\mathcal{SW}(\frac{3}{2}, d)$ -algebras. All symbolic calculations have been performed in MATHEMATICATM and REDUCE while for the time consuming commutation operations and the expansion of NOP's a special C-program had to be used.

Since the values of the coupling constants do not depend on the special choice of the basis in the space of quasiprimary fields we will in general omit the coupling constants involving non-simple fields. A special choice of a quasiprimary basis is contained in appendix A and B. In appendix C we list the Kac-determinants of states involving exactly one additional field.

The cases $\frac{3}{2} \leq d \leq \frac{7}{2}$ were treated by J.Figueroa-O'Farrill et al. [17].

$\mathcal{SW}(\frac{3}{2}, \frac{3}{2})$

This algebra is a super Lie algebra with the followig commutation relations:

$$\begin{aligned}
[\phi_m, \phi_n]_+ &= C_{\phi\phi}^{\psi} \psi_{m+n} + 2L_{m+n} + \frac{c}{3}(m^2 - \frac{1}{4})\delta_{m+n,0} \\
[\phi_m, \psi_n] &= \frac{1}{3}(n - 2m)(C_{\phi\psi}^{\phi} \phi_{m+n} + \frac{3}{2}G_{m+n}) \\
[\psi_m, \psi_n] &= \frac{n - m}{2}C_{\psi\psi}^{\psi} \psi_{m+n} + (n - m)L_{m+n} + \frac{c}{12}(n^3 - n)\delta_{m+n,0}
\end{aligned}$$

where $C_{\phi\phi}^{\psi} = C_{\psi\psi}^{\psi}$ and $C_{\phi\psi}^{\phi} = \frac{3}{4}C_{\psi\psi}^{\psi}$

Thus $\mathcal{SW}(\frac{3}{2}, \frac{3}{2})$ exists generically and the coupling constant $C_{\psi\psi}^\psi$ is a free parameter. Note that one may choose a new basis, such that the resulting commutators define the direct sum of two super Virasoro algebras. This case was discussed in [17] and is similar to the conformal one treated by Zamolodchikov [2].

The cases $d = 2, \frac{5}{2}$ with vanishing self-coupling have been investigated in detail by T. Inami et al. [14].

$\mathcal{SW}(\frac{3}{2}, 2)$

This is the first case where a $\mathcal{SW}$ -algebra does not close linearly. The commutation relations are given by:

$$\begin{aligned}
[\phi_m, \phi_n] &= \frac{n-m}{2} C_{\phi\phi}^\phi \phi_{m+n} + (n-m) L_{m+n} + \frac{c}{12} (n^3 - n) \delta_{m+n,0} \\
[\phi_m, \psi_n] &= p_{2, \frac{5}{2}, \frac{7}{2}}(m, n) (C_{\phi\psi}^{\mathcal{N}(\phi, G)} \mathcal{N}(\phi, G)_{m+n} + C_{\phi\psi}^{\mathcal{N}(L, G)} \mathcal{N}(L, G)_{m+n}) + \\
&\quad p_{2, \frac{5}{2}, \frac{5}{2}}(m, n) C_{\phi\psi}^\psi \psi_{m+n} + \\
&\quad p_{2, \frac{5}{2}, \frac{3}{2}}(m, n) \frac{3}{5} C_{G\phi}^\psi G_{m+n} \\
[\psi_m, \psi_n]_+ &= p_{\frac{5}{2}, \frac{5}{2}, 4}(m, n) (C_{\psi\psi}^{\mathcal{N}(\phi, \phi)} \mathcal{N}(\phi, \phi)_{m+n} + C_{\psi\psi}^{\mathcal{N}(\psi, G)} \mathcal{N}(\psi, G)_{m+n} + \\
&\quad C_{\psi\psi}^{\mathcal{N}(\phi, L)} \mathcal{N}(\phi, L)_{m+n} + C_{\psi\psi}^{\mathcal{N}(G, \partial G)} \mathcal{N}(G, \partial G)_{m+n} + \\
&\quad C_{\psi\psi}^{\mathcal{N}(L, L)} \mathcal{N}(L, L)_{m+n}) + \\
&\quad p_{\frac{5}{2}, \frac{5}{2}, 2}(m, n) (C_{\psi\psi}^\phi \phi_{m+n} + 2L_{m+n}) + \\
&\quad \frac{2c}{5} \binom{n + \frac{3}{2}}{4} \delta_{m+n,0}
\end{aligned}$$

where $C_{\psi\psi}^\phi = \frac{2}{5} C_{\phi\phi}^\phi$, $C_{\phi\psi}^\psi = \frac{1}{2} C_{\phi\phi}^\phi$ and

$$(C_{\phi\phi}^\phi)^2 = \frac{4(5c+6)^2}{(4c+21)(15-c)}.$$

We have verified explicitly that the coupling constant $C_{\psi\psi}^{\mathcal{N}(\phi, \phi)}$ is zero. In general, the field $\mathcal{N}(\phi, \phi)$ cannot be neglected since it contributes to the d -matrix. Only for $c = -\frac{6}{5}$ the self-coupling vanishes and all fields containing ϕ or ψ can be ignored. In this case the fields $\mathcal{N}(\phi, \phi)$ and $\mathcal{N}(\psi, \phi)$ have vanishing primary projection. Generically, this algebra contains a super primary field of dimension 4 which is quadratic in the simple super primary field. We have verified that if this field occurs in a Jacobi identity the factor in front of it vanishes (cf. remark at the end of chapter 3).

Thus the algebra exists generically, which is not surprising since the classical counterpart is the symmetry algebra of the super Toda theory corresponding to the super Lie algebra $Osp(3|2)$.

$\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$

The self-coupling vanishes and only two c values are possible:

$$c = -\frac{5}{2}, \frac{10}{7}$$

$\mathcal{SW}(\frac{3}{2}, 3)$

Here the self-coupling is zero again and the allowed c values are:

$$c = -\frac{45}{2}, -\frac{27}{7}, \frac{5}{4}$$

$\mathcal{SW}(\frac{3}{2}, \frac{7}{2})$

This is a case with non-vanishing self-coupling and consistency is obtained for:

c	$(C_{\Phi\Phi}^\Phi)^2$
$\frac{7}{5}$	$\frac{146016}{79135}$
$-\frac{17}{11}$	$-\frac{45009216}{523393}$

In the sequel we present new $\mathcal{SW}(\frac{3}{2}, d)$ -algebras with d up to 7.

$\mathcal{SW}(\frac{3}{2}, 4)$

While for $c = -\frac{20}{3}$ the algebra $\mathcal{SW}(\frac{3}{2}, 4)$ is consistent for vanishing self-coupling, it is also consistent with nonzero self-coupling for four values of c :

c	$(C_{\Phi\Phi}^\Phi)^2$
$-\frac{185}{4}$	$\frac{355359375}{161644468}$
-13	$\frac{427063}{4247}$
$-\frac{21}{2}$	$-\frac{508369}{2499}$
$-\frac{120}{13}$	$-\frac{273043750000}{7804122111}$
$-\frac{20}{3}$	0

Only in the case $c = -\frac{120}{13}$ the field $\mathcal{N}(\phi, \phi)$ has a nontrivial primary projection and even in this case the coupling constant $C_{\psi\psi}^{\mathcal{N}(\phi, \phi)}$ vanishes. This means that there is a primary field of dimension 8 contained in the algebra at $c = -\frac{120}{13}$. In this case the mentioned field contributes to the d matrix.

$\mathcal{SW}(\frac{3}{2}, \frac{9}{2})$

This algebra exists only with vanishing self-coupling for three values of the central charge:

$$c = -\frac{69}{2}, -\frac{81}{10}, \frac{4}{11}$$

$\mathcal{SW}(\frac{3}{2}, 5)$

Analogously to $\mathcal{SW}(\frac{3}{2}, 3)$ the self-coupling has to vanish. Consistency implies here $c = -\frac{105}{11}$.

$\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$

The algebra $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ is consistent for five values of c :

c	$(C_{\Phi\Phi}^\Phi)^2$
$\frac{10}{7}$	$\frac{844918800}{34554863}$
$\frac{11}{40}$	$-\frac{91376712973}{3514069779}$
$-\frac{155}{19}$	$\frac{394077798400}{181920009}$
$-\frac{705}{8}$	$-\frac{4322235328125}{3075066706603}$
$-\frac{5}{13}$	0

$\mathcal{SW}(\frac{3}{2}, 6)$

This algebra is consistent with non-vanishing self-coupling for

c	$(C_{\Phi\Phi}^\Phi)^2$
$-\frac{33}{2}$	$\frac{6309688448}{3137409}$
-18	$-\frac{16916841216}{17064005}$
$-\frac{2241}{20}$	$\frac{788850108780331059}{143584629321555025}$

With vanishing self-coupling it exists for:

$$c = -\frac{93}{2}, -\frac{162}{13}, \frac{27}{20}$$

The field $\mathcal{N}(\phi, \phi)$ has a non-vanishing primary projection only for $c = -18$ and $-\frac{2241}{20}$. In these cases $C_{\psi\psi}^{\mathcal{N}(\phi, \phi)}$ is zero.

$\mathcal{SW}(\frac{3}{2}, \frac{13}{2})$

This algebra exists only for zero self-coupling and $c = -\frac{195}{14}$.

$\mathcal{SW}(\frac{3}{2}, 7)$

Finally we discuss the algebra obtained by adding a super primary field of dimension 7. Since 7 is odd the self-coupling vanishes. For this algebra we have checked the condition $(\phi_m, \phi_n, \phi_k, \phi_{m+n+k}) = 0$ only. The only allowed c values are:

$$c = -\frac{77}{5}, -\frac{13}{8}$$

In the next section we will give some arguments that for these values of the central charge the algebra should indeed exist.

5. Structure of $\mathcal{SW}(\frac{3}{2}, d)$ -algebras

In this chapter we summarize our results concerning $\mathcal{SW}$ -algebras with two generators. We will show that in perfect analogy to $\mathcal{W}(2, \delta)$ -algebras for most values of the central charge a classification into special series is possible.

$\mathcal{W}(2, \delta)$ -algebras can be classified by the values of the central charge. Until now five different classes appeared:

- the generically existing algebras related to simple Lie algebras (see e.g. [23]),
- the algebras related to the ADE classification of A. Cappelli et al. [24][10],
- the algebras of the (1,s) series [25],
- the parabolic $\mathcal{W}(2, \delta)$ -algebras related to degenerate representations of the Virasoro algebra [26] and
- some exceptional cases [10][27].

For $\mathcal{SW}(2, d)$ -algebras these five types also exist as will be shown below.

The algebras $\mathcal{SW}(\frac{3}{2}, d)$ with $\frac{3}{2} \leq d \leq 7$ exist for the following values of the central charge:

d	c
$\frac{3}{2}$	generic
2	generic
$\frac{5}{2}$	$-\frac{5}{2}, \frac{10}{7}$
3	$-\frac{45}{2}, -\frac{27}{7}, \frac{5}{4}$
$\frac{7}{2}$	$-\frac{17}{11}, \frac{7}{5}$
4	$-\frac{185}{4}, -13, -\frac{21}{2}, -\frac{120}{13}, -\frac{20}{3}$
$\frac{9}{2}$	$-\frac{69}{2}, -\frac{81}{10}, \frac{4}{11}$
5	$-\frac{105}{11}$
$\frac{11}{2}$	$-\frac{705}{8}, -\frac{155}{19}, -\frac{5}{13}, \frac{11}{40}, \frac{10}{7}$
6	$-\frac{2241}{20}, -\frac{93}{2}, -18, -\frac{33}{2}, -\frac{162}{13}, \frac{27}{20}$
$\frac{13}{2}$	$-\frac{195}{14}$
7	$-\frac{77}{5}, -\frac{13}{8}$

First, the generically existing algebras $\mathcal{SW}(\frac{3}{2}, \frac{3}{2})$ and $\mathcal{SW}(\frac{3}{2}, 2)$ are closely related to the Lie super algebras $Osp(2|1)$ and $Osp(3|2)$, respectively [14].

Secondly, all super minimal c values can be related to the ADE classification of modular invariant partition functions of A. Cappelli et al. [24]. To be more specific, we need the

minimal models of the super Virasoro algebra. Therefore we list the values of the central charge and the corresponding superconformal dimensions [28][29][30][31].

$$\begin{aligned}
c(p, q) &= \frac{3}{2} \left(1 - \frac{2(p-q)^2}{pq} \right) \quad \text{with either } p, q \in \mathbb{N}, p, q \text{ coprime and } p+q \in 2\mathbb{N} \\
&\quad \text{or} \quad p, q \in 2\mathbb{N}, \frac{p}{2}, \frac{q}{2} \text{ coprime and } \frac{p}{2} + \frac{q}{2} \notin 2\mathbb{N} \\
h(r, s) &= \frac{(rp - qs)^2 - (p - q)^2}{8pq} + \frac{1 - (-1)^{r+s}}{32} \quad 1 \leq r \leq q-1, 1 \leq s \leq p-1
\end{aligned} \tag{5.1}$$

$r + s$ even yields representations in the Neveu-Schwarz sector and $r + s$ odd those in the Ramond sector.

The divisibility conditions in (5.1) ensure that $p, q \in \mathbb{N}$ with $p + q \in 2\mathbb{N}$ are chosen as small as possible.

Some simple fusion rule arguments suggest the following

Proposition: For any $d \in \mathbb{N}$ with $d = \frac{1}{8}(p-2)(q-2)$, $p, q \in 2\mathbb{N}$, $\frac{p}{2}, \frac{q}{2}$ coprime and $\frac{p+q}{2} \in 2\mathbb{N}+1$ the algebra $\mathcal{SW}(\frac{3}{2}, d)$ exists for $c = c(p, q)$ with vanishing self-coupling.

Since in this case the central charge belongs to the super minimal series and the dimension of the additional super primary field Φ can be parametrized by $d = h(1, p-1) = h(q-1, 1)$, the well known fusion rules for super Virasoro minimal models may be applied [28][29][30]. In terms of superconformal families they read

$$[\Phi] \times [\Phi] = [1]$$

such that the additional field of dimension d is a ‘simple current’ in the sense of [32]. In this case general arguments suggest that this ‘simple current’ can be added to the algebra of the superconformal family of the identity, although this has been proven only for the unitary case [32][33][34][35].

Many examples with zero self-coupling may be explained by the proposition. In fact the two c values for which $\mathcal{SW}(\frac{3}{2}, 7)$ might exist belong to this series, so one may assume that $\mathcal{SW}(\frac{3}{2}, 7)$ indeed exists for these two values of the central charge.

By considering the chiral part of the explicit form of a modular invariant partition function an interpretation for all super minimal values of the central charge is possible [24]:

series	$c(p, q) = (p, q)$	$h(r, s) = (r, s)$
$(D_{2\rho+2}, A_{q-1})$	$(4\rho + 2, q)$	$(q - 1, 1) = \frac{1}{2}\rho(q - 2)$
$(E_6, A_{\rho-2})$	$(\rho - 1, 12)$	$(7, 1) = \frac{\rho-4}{2}$
$(E_6, D_{\rho+1})$	$(2\rho, 12)$	$(5, 1) = \frac{\rho-2}{2}$ $(7, 1) = \frac{2\rho-3}{2}$ $(11, 1) = \frac{5(\rho-1)}{2}$
$(E_8, A_{\rho-2})$	$(\rho - 1, 30)$	$(11, 1) = \frac{\rho-6}{2}$ $(19, 1) = \frac{3(\rho-4)}{2}$ $(29, 1) = \frac{7(\rho-3)}{2}$

Note that all fields corresponding to the dimensions $h(r, s)$ given above are local.

Because the partition functions related to the first two rows of the table consist of binomials only, the number of simple fields in corresponding RCFTs is at most two. This case is closely related to $\mathcal{SW}$ -algebras with two generators. Indeed the $(D_{2\rho+2}, A_{q-1})$ series leads exactly to the cases explained by our proposition. Furthermore, note that in the $(E_6, D_{\rho+1})$ and the $(E_8, A_{\rho-2})$ series the number of simple fields is at most 4. So there seems to be a connection to $\mathcal{SW}$ -algebras with four generators. Taking into account that the commutators of the super primary fields corresponding to the first h values cannot contain the other fields leads to series of $\mathcal{SW}(\frac{3}{2}, d)$ -algebras, too. For consistency the other super primary fields contained in these theories cannot be simple.

There are two examples that fit in the $(E_6, A_{\rho-2})$ series both with $C_{\Phi\Phi}^\Phi \neq 0$. Examples are $\mathcal{SW}(\frac{3}{2}, \frac{7}{2})$ at $c = \frac{7}{5}$ and $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ at $\frac{10}{7}$. This series is realized only for odd ρ leading to $(p, q) = (\rho - 1, 12)$ that fulfill the divisibility condition in (5.1). These two Algebras can also be related to the $(E_6, D_{\rho+1})$ series where the dimension d is given by the second h value in the table. The $(E_6, D_{\rho+1})$ series is realized for $\rho = 7, 11, 13$ as $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$, $\mathcal{SW}(\frac{3}{2}, \frac{9}{2})$ and $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ with vanishing self-coupling. Note that only the cases where ρ and 6 are coprime appear. Since $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$ and $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ exist for $c = \frac{10}{7}$ and both can be related to (E_6, D_8) there should be a connection. We have explicitly verified that $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$ contains a super primary non-simple field of dimension $\frac{11}{2}$ at $c = \frac{10}{7}$. There is strong evidence that this yields a realization of $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ at the c value mentioned because we have calculated the self-coupling of one component of the non-simple super primary field. Its value is exactly that obtained by the direct construction of $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$. For details we refer to appendix D. A similar construction where $\mathcal{SW}(\frac{3}{2}, \frac{7}{2})$ is realized as a subalgebra of $\mathcal{SW}(\frac{3}{2}, \frac{3}{2})$ at $c = \frac{7}{5}$ has been carried out by K. Hornfeck [19].

The $(E_8, A_{\rho-2})$ series is realized only for $\rho = 17$ as $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ at $c = \frac{11}{40}$.

For $12 \leq \rho \leq 16$ the resulting pair $(\rho - 1, 30)$ does not belong to the $(E_8, A_{\rho-1})$ series since the divisibility condition in (5.1) is not satisfied. The next possible value for ρ is 23 and the corresponding dimension $\frac{17}{2}$.

Thirdly, note that $\mathcal{SW}(\frac{3}{2}, d)$ exists for $c = c(1, s)$, s odd and $d = \frac{2s-1}{2}$. This case is similar to the conformal one treated by H.G. Kausch for $\mathcal{W}$ -algebras with two generators. In [25] a free field construction was presented and one can hope that similar techniques lead to a free field realization in the supersymmetric case. We will call this series the $(1, s)$ -series.

Fourthly, all $\mathcal{SW}(\frac{3}{2}, d)$ with $d = \frac{3}{2}n$ or $d = 2n$, $n \in \mathbb{N}$ exist for $c = \frac{3}{2}(1 - \frac{16}{3}d) = \frac{3}{2}(1 - 8n) = c_d^3(n)$ or $c = \frac{3}{2}(1 - 2d) = \frac{3}{2}(1 - 4n) = c_d^8(n)$ respectively. These cases are related to degenerate representations of the super Virasoro algebra and will be called parabolic. The central charge and the dimensions of the primary fields for the super Virasoro algebra are given by [29]:

$$\begin{aligned} c &= \frac{3}{2}(1 - 16\alpha_0^2) \\ \alpha_{\pm} &= \sqrt{\alpha_0^2 + \frac{1}{2}} \pm \alpha_0 \\ h_{r,s} &= \frac{(r\alpha_+ - s\alpha_-)^2}{4} - \alpha_0^2 \end{aligned} \tag{5.2}$$

Note that $h_{r,r} = \alpha_0^2(r^2 - 1)$ and $h_{r,-r} = \alpha_0^2(r^2 - 1) + \frac{1}{2}r^2$. The series described above can be obtained by adding an additional field of superconformal dimension $d = h_{2,2}$ or $d = h_{3,3}$ to the superconformal family of the identity. Note that these fields are local because they obey the locality condition

$$\epsilon_{dd} = e^{2\pi i(1-r)^2\alpha_0^2} = e^{2\pi i\frac{(1-r)^2}{r^2-1}d} = \pm 1$$

if one assumes $2\frac{(r-1)^2}{r^2-1}d \in \mathbb{N}$. Taking into account that the fields corresponding to $h_{2,2}$ or $h_{3,3}$ are either bosonic or fermionic leads to the possible values of d given above. For $r \geq 4$ one would not get a $\mathcal{SW}$ -algebra with only two generators. These algebras lead to rational theories with effective central charge $\tilde{c} = \frac{3}{2}$ [36]. For the similar case of $\mathcal{W}(2, \delta)$ -algebras the characters and S -matrices have been calculated and the generalization to the super case is straightforward [26].

Finally, some algebras remain which do not belong to any of the series described above. All of them have non-vanishing self-coupling. The investigation of the representation theory of these algebras leads to a better understanding. One obtains e.g. that the effective central charge is greater than $\frac{3}{2}$ [36].

Finally we list all $\mathcal{SW}(\frac{3}{2}, d)$ -algebras with $\frac{3}{2} \leq d \leq 7$ and show how they fit into these patterns.

d	$c = c(p, q)$		$d = h(r, s)$	series
	$C_{\Phi\Phi}^\Phi = 0$	$C_{\Phi\Phi}^\Phi \neq 0$		
$\frac{3}{2}$	generic	generic		
2	$(10, 4) = -\frac{6}{5}$	generic	(3,1)	(D_6, A_3)
$\frac{5}{2}$	$(3, 1) = -\frac{5}{2}$ $(14, 12) = \frac{10}{7}$		(3,1) (5,1)	(1, s) (E_6, D_8)
3	$(14, 4) = -\frac{27}{7}$ $(6, 8) = \frac{5}{4}$ $c_d^3(2) = -\frac{45}{2}$		(3,1) (7,1) (2,2)	(D_8, A_3) (D_4, A_7) parabolic
$\frac{7}{2}$		$(10, 12) = \frac{7}{5}$ $-\frac{17}{11}$	(7,1)	(E_6, A_9) ?
4	$(18, 4) = -\frac{20}{3}$	$c_d^8(2) = -\frac{21}{2}$ $-\frac{185}{4}$ -13 $-\frac{120}{13}$	(3,1) (3,3)	(D_{10}, A_3) parabolic ? ? ?
$\frac{9}{2}$	$(5, 1) = -\frac{81}{10}$ $(22, 12) = \frac{4}{11}$ $c_d^3(3) = -\frac{69}{2}$		(3,1) (5,1) (2,2)	(1, s) (E_6, D_{12}) parabolic
5	$(22, 4) = -\frac{105}{11}$		(3,1)	(D_{12}, A_3)
$\frac{11}{2}$	$(26, 12) = -\frac{5}{13}$	$(14, 12) = \frac{10}{7}$ $(16, 30) = \frac{11}{40}$ $-\frac{705}{8}$ $-\frac{155}{19}$	(5,1) (7,1) (11,1)	(E_6, D_{14}) (E_6, A_{13}) (E_8, A_{15}) ? ?
6	$(26, 4) = -\frac{162}{13}$ $(10, 8) = \frac{27}{20}$ $c_d^3(4) = -\frac{93}{2}$	$c_d^8(3) = -\frac{33}{2}$ -18 $-\frac{2241}{20}$	(3,1) (7,1) (2,2) (3,3)	(D_{14}, A_3) (D_6, A_7) parabolic parabolic ? ?
$\frac{13}{2}$	$(7, 1) = -\frac{195}{14}$		(3,1)	(1, s)
7	$(30, 4) = -\frac{77}{5}$ $(6, 16) = -\frac{13}{8}$		(3,1) (15,1)	(D_{16}, A_3) (D_4, A_{15})

6. Explicit results about $\mathcal{SW}(\frac{3}{2}, \mathbf{d}_1, \mathbf{d}_2)$ -algebras

In this chapter we present our results about the construction of $\mathcal{SW}$ -algebras with two additional generators. For these algebras one has to check the validity of 20 Jacobi identities. For $\mathcal{SW}(\frac{3}{2}, d_1, d_2)$ we will denote the two additional super fields of dimension d_i by $\Phi^i = \phi^i + \theta\psi^i$.

$\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, \frac{3}{2})$

In this case there are five free parameters namely the central charge and four coupling constants. Consistency implies that the commutators close linearly in the fields and also the following condition:

$$(C_{\phi^1\phi^1}^{\psi^2})^2 + (C_{\phi^2\phi^2}^{\psi^1})^2 - C_{\phi^1\phi^1}^{\psi^2}C_{\phi^2\phi^2}^{\psi^2} - C_{\phi^2\phi^2}^{\psi^1}C_{\phi^1\phi^1}^{\psi^1} - 4 = 0$$

In complete analogy to the $\mathcal{SW}(\frac{3}{2}, \frac{3}{2})$ -algebra one can choose a new basis such that the resulting commutators define the direct sum of three super Virasoro algebras. As described in [17] this case can be generalized easily to the extension of the super Virasoro algebra by n simple fields of dimension $\frac{3}{2}$.

The following algebra has been studied in [18] using the covariant approach.

$\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, 2)$

For this algebra five parameters have to be calculated. The Jacobi identities imply that two sets of solutions exist:

1.)

$$\begin{aligned} C_{\phi^1\phi^1}^{\phi^2} &= 0 \\ C_{\phi^1\phi^1}^{\psi^1} &= \frac{(C_{\phi^2\phi^2}^{\psi^1})^2 - 4}{C_{\phi^2\phi^2}^{\psi^1}} \\ (C_{\phi^2\phi^2}^{\phi^2})^2 &= \frac{4(12 + 10c + 3(C_{\phi^2\phi^2}^{\psi^1})^2)^2 (4 + (C_{\phi^2\phi^2}^{\psi^1})^2)}{(84 + 16c + 21(C_{\phi^2\phi^2}^{\psi^1})^2)(60 - 4c + 15(C_{\phi^2\phi^2}^{\psi^1})^2)} \end{aligned}$$

2.)

$$\begin{aligned} C_{\phi^2\phi^2}^{\psi^1} &= \frac{1}{2}C_{\phi^1\phi^1}^{\psi^1} \\ (C_{\phi^1\phi^1}^{\psi^1})^2 &= 4(C_{\phi^1\phi^1}^{\phi^2})^2 - 4C_{\phi^1\phi^1}^{\phi^2}C_{\phi^2\phi^2}^{\phi^2} - 16 \\ C_{\phi^1\phi^1}^{\phi^2} &= \frac{8c}{10c - 27}C_{\phi^2\phi^2}^{\phi^2} \end{aligned}$$

This means that in both cases there remain two free parameters. While the first solution is a trivial one as discussed below the second solution could be the symmetry algebra of the quantized Toda theory corresponding to $D(2|1, \alpha)$.

$\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, \frac{5}{2})$

Since the coupling constant of the superconformal family of dimension $\frac{3}{2}$ to the superconformal family of dimension $\frac{5}{2}$ and the self-coupling of the latter are zero there remain three free parameters. Two types of solutions exist with vanishing $C_{\phi^1\phi^1}^{\psi^2}$:

1.)

$$\begin{aligned} (C_{\phi^2\phi^2}^{\psi^1})^2 &= \frac{14c-20}{5} & C_{\phi^1\phi^1}^{\psi^1} C_{\phi^2\phi^2}^{\psi^1} &= \frac{2(7c-20)}{5} & c &\neq \frac{10}{7} \\ \text{or} \quad (C_{\phi^2\phi^2}^{\psi^1})^2 &= -\frac{8c+20}{5} & C_{\phi^1\phi^1}^{\psi^1} C_{\phi^2\phi^2}^{\psi^1} &= -\frac{8(c+5)}{5} & c &\neq -\frac{5}{2} \end{aligned}$$

2.)

$$(C_{\phi^2\phi^2}^{\psi^1})^2 = -4 \quad C_{\phi^1\phi^1}^{\psi^1} C_{\phi^2\phi^2}^{\psi^1} = -8$$

In both cases the central charge is free.

For $\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, 2)$ and $\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, \frac{5}{2})$ the first solutions can be decomposed into a direct sum. In general, one has the following structure. Assume that $\mathcal{SW}(\frac{3}{2}, d)$ with generators $\hat{\mathcal{L}}$ and $\hat{\Phi}$ exists for $\hat{c}$ and self-coupling $\hat{C}_{\hat{\Phi}\hat{\Phi}}^{\hat{\Phi}} = f(\hat{c})$. Then $\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, d)$ exists for generic central charge c and:

$$\begin{aligned} C_{\phi^1\phi^1}^{\psi^2} &= 0 & (C_{\phi^1\phi^1}^{\psi^1})^2 &= \frac{4}{\hat{c}} \frac{(c-2\hat{c})^2}{c-\hat{c}} \\ (C_{\phi^2\phi^2}^{\psi^1})^2 &= 4 \frac{c-\hat{c}}{\hat{c}} & C_{\Phi\Phi}^{\Phi} &= \frac{c}{\hat{c}} f(4(4+C_{\phi^2\phi^2}^{\psi^1})^{-1}) \end{aligned}$$

This is easily seen since a change of basis implies for this solution:

$$\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, d) \cong \mathcal{SW}(\frac{3}{2}) \oplus \mathcal{SW}(\frac{3}{2}, d)$$

For the second solution of $\mathcal{SW}(\frac{3}{2}, \frac{3}{2}, \frac{5}{2})$ such a change of basis becomes singular and is therefore not possible. It is remarkable that one can choose a linear combination of the super Virasoro field and the additional super field of dimension $\frac{3}{2}$ such that the resulting field (anti-)commutes with itself. This field is built up by the sum of the Super Virasoro field and $\frac{C_{\phi^1\phi^1}^{\psi^1}}{4}$ times the field Φ^1 .

$\mathcal{SW}(\frac{3}{2}, 2, 2)$

Consistency implies fixed central charge and the following conditions for the four free coupling constants:

$$\begin{aligned} c &= \frac{3}{2} \\ C_{\phi^2\phi^2}^{\phi^1} &= -C_{\phi^1\phi^1}^{\phi^1} & C_{\phi^1\phi^1}^{\phi^2} &= -C_{\phi^2\phi^2}^{\phi^2} \\ (C_{\phi^1\phi^1}^{\phi^1})^2 &+ (C_{\phi^2\phi^2}^{\phi^2})^2 &= 2 \end{aligned}$$

For this solution the fields $\mathcal{N}(\phi^1, \phi^1)$, $\mathcal{N}(\phi^2, \phi^2)$ and $\mathcal{N}(\phi^1, \phi^2)$ have no primary projection and do not contribute to the algebra.

Obviously, this algebra has an inner $SO(2)$ symmetry realized as a rotation in the space of the super fields of dimension two. Consequently the solution is determined by relations that are invariant under the action of $SO(2)$. Note that a rotation by an angle α in the space of super fields yields a rotation of angle 3α in the space of self-couplings.

One could speculate that $\mathcal{SW}(\frac{3}{2}, 2, 2)$ for $c = \frac{3}{2}$ is a subalgebra of $\mathcal{SW}(\frac{3}{2}, 2, 2, \frac{7}{2})$, which should exist generically and is related to $Osp(4|4)$ [37]. A very similar case for the $\mathcal{W}(2, 4, 4)$ -algebra which is related to $SO(8)$ has been treated in [25].

$\mathcal{SW}(\frac{3}{2}, 2, \frac{5}{2})$

For this algebra there exist two consistent sets of solutions:

1.)

$$\begin{aligned} c &= -15 \\ (C_{\phi^2\phi^2}^{\phi^1})^2 &= -\frac{40}{13} \\ (C_{\phi^1\phi^1}^{\phi^1})^2 &= -\frac{1058}{65} \\ C_{\phi^2\phi^2}^{\phi^1} C_{\phi^1\phi^1}^{\phi^1} &= -\frac{92}{13} \end{aligned}$$

2.)

$$\begin{aligned} c &= \frac{39}{2} \\ (C_{\phi^2\phi^2}^{\phi^1})^2 &= -88 \\ (C_{\phi^1\phi^1}^{\phi^1})^2 &= -\frac{1058}{11} \\ C_{\phi^2\phi^2}^{\phi^1} C_{\phi^1\phi^1}^{\phi^1} &= -92 \end{aligned}$$

For $c = -15$ the field $\mathcal{N}(\phi^1, \phi^2)$ can be written as a linear combination of the other fields of dimension $\frac{9}{2}$. This is the only case in which null fields appear and must be omitted (cf. the remark at the end of chapter 3).

$\mathcal{SW}(\frac{3}{2}, \frac{5}{2}, \frac{5}{2})$

There is no solution for this algebra. Analogously to $\mathcal{W}$ -algebras a $\mathcal{SW}$ -algebra with two additional super fields and no nonzero coupling constant cannot exist, since a Jacobi identity of type $([[\phi^1_m, \phi^1_n]_{\pm}, \phi^2_k]_{\pm})_{cycl.} = 0$ cannot be satisfied [10].

That means $\mathcal{SW}(\frac{3}{2}, d_1, d_2)$ cannot exist for $[d_1 + \frac{1}{2}], [d_2 + \frac{1}{2}] \in 2\mathbb{N} + 1$.

$\mathcal{SW}(\frac{3}{2}, \frac{5}{2}, \frac{7}{2})$

In this case two coupling constants are free and one obtains only one set of solutions with

no free parameters:

$$\begin{aligned}
c &= \frac{13}{6} \\
(C_{\phi^1\phi^1}^{\psi^2})^2 &= \frac{5208}{979} \\
(C_{\phi^2\phi^2}^{\psi^2})^2 &= \frac{317709}{60698} \\
C_{\phi^1\phi^1}^{\psi^2} C_{\phi^2\phi^2}^{\psi^2} &= -\frac{5166}{979}
\end{aligned}$$

This is an algebra predicted by Schoutens et al. in [38] by coset considerations. It has been shown in [19] by K. Hornfeck that if this $\mathcal{SW}$ -algebra has a $\mathcal{W}(2, 3, 4)$ as subalgebra it can at most exist for $c = \frac{13}{6}$. Indeed this value is the only possible one.

7. Conclusion

Using a non-explicitly covariant approach we have been able to construct a whole bunch of new $\mathcal{SW}$ -algebras with two generators. Most of the results fit into systematic patterns in complete analogy to $\mathcal{W}$ -algebras. For some new $\mathcal{SW}$ -algebras namely $\mathcal{SW}(\frac{3}{2}, 4)$ at $c = -\frac{185}{4}, -\frac{120}{13}, -13$ and $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ at $c = -\frac{705}{8}, -\frac{155}{19}$ and $\mathcal{SW}(\frac{3}{2}, 6)$ at $c = -18, -\frac{2241}{20}$ an interpretation has not been found yet. Even the interpretation of $\mathcal{SW}(\frac{3}{2}, \frac{7}{2})$ at $c = -\frac{17}{11}$ is still an open question although this algebra has been known for some time. A study of the possible and physically relevant highest weight representations of these algebras will lead to a better understanding of these algebras. The methods explained in [39][27] can be applied to this case with only small changes and work is in progress [36].

In contrast to $\mathcal{W}(2, \delta)$ -algebras no consistent $\mathcal{SW}(\frac{3}{2}, d)$ -algebras with irrational values of the central charge appeared. Furthermore, there is only one generically existing nonlinear $\mathcal{SW}(\frac{3}{2}, d)$ -algebra, namely $\mathcal{SW}(\frac{3}{2}, 2)$.

For $\mathcal{SW}$ -algebras with three generators we have constructed two new algebras that exist for finitely many c values only.

Since the difficulties of the transition from $\mathcal{W}$ -algebras to $\mathcal{SW}$ -algebras have not been too hard we are confident that similar calculations for the super $N = 2$ case will be possible in the near future.

Acknowledgements

We would like to thank W. Nahm, M. Flohr, J. Kellendonk, S. Mallwitz, A. Recknagel, M. Rösger, M. Terhoeven and R. Varnhagen for many useful discussions. The creative atmosphere in the institute was an important support for this work.

It is a pleasure to thank the Max-Planck-Institut für Mathematik in Bonn-Beuel especially Th. Berger, S. Mauermann and T. Höfer since nearly all calculations have been performed in their computer centre.

Appendix:

In the appendices A, B and C we list all fields and determinants (of matrices (d_{ij})) which occur in the examples considered in chapter 4. Because fields of different conformal dimensions are orthogonal, the matrices have block-diagonal form. Furthermore, all fields in the superconformal family of the identity have vanishing (d_{ij}) with fields of the superconformal family of the additional super primary field. Thus the determinants factorize in two terms:

$$\det D_\delta = \det D_\delta^{[1]} \det D_\delta^{[d]}$$

where $[d]$ stands for the superconformal family of the additional super primary field of dimension d and δ is the dimension considered. But there is one exception to this rule: if $d \in 2\mathbb{N}$ and $\delta = 2d$ the field $\mathcal{N}(\phi, \phi)$ is involved and the determinant $\det D_\delta$ does not factorize. Because such determinants are very complicated we omit them.

Appendix A: A basis of quasiprimary fields up to dimension $\frac{25}{2}$ built up by $\mathcal{L}$

$$d = \frac{3}{2} : 1 \text{ field} : G$$

$$\det D_{\frac{3}{2}}^{[1]} = \frac{2c}{3}$$

$$d = 2 : 1 \text{ field} : L$$

$$\det D_2^{[1]} = \frac{c}{2}$$

$$d = \frac{7}{2} : 1 \text{ field} : \mathcal{N}(L, G)$$

$$\det D_{\frac{7}{2}}^{[1]} = \frac{1}{12}c(21 + 4c)$$

$$d = 4 : 2 \text{ fields} : \mathcal{N}(G, \partial G), \mathcal{N}(L, L)$$

$$\det D_4^{[1]} = \frac{1}{60}c^2(21 + 4c)(-7 + 10c)$$

$$d = \frac{9}{2} : 1 \text{ field} : \mathcal{N}(L, \partial G)$$

$$\det D_{\frac{9}{2}}^{[1]} = -\frac{2}{35}c(-7 + 10c)$$

$$d = \frac{11}{2} : 2 \text{ fields} : \mathcal{N}(L, \partial^2 G), \mathcal{N}(\mathcal{N}(L, L), G)$$

$$\det D_{\frac{11}{2}}^{[1]} = \frac{1}{54}c^2(11 + c)(21 + 4c)(-7 + 10c)$$

$$d = 6 : 4 \text{ fields} :$$

$$\mathcal{N}(G, \partial^3 G), \mathcal{N}(\mathcal{N}(L, G), \partial G), \mathcal{N}(\mathcal{N}(L, L), L), \mathcal{N}(L, \partial^2 L)$$

$$\det D_6^{[1]} = \frac{1}{378}c^4(11 + c)(21 + 4c)^2(-7 + 10c)^2(11 + 14c)$$

$$d = \frac{13}{2} : 2 \text{ fields} :$$

$$\mathcal{N}(L, \partial^3 G), \mathcal{N}(\mathcal{N}(L, L), \partial G)$$

$$\det D_{\frac{13}{2}}^{[1]} = \frac{16}{847}c^2(21 + 4c)(-7 + 10c)(11 + 14c)$$

$$d = 7 : 1 \text{ field} : \mathcal{N}(\mathcal{N}(L, G), \partial^2 G)$$

$$\det D_7^{[1]} = -\frac{2}{45}c(21 + 4c)(-7 + 10c)$$

$d = \frac{15}{2} : 5$ fields :

$$\mathcal{N}(\mathcal{N}(G, \partial G), \partial^2 G), \quad \mathcal{N}(L, \partial^4 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^2 G), \\ \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 L), G)$$

$$\det D_{\frac{15}{2}}^{[1]} = \frac{64}{27885}(-1+c)c^5(11+c)(21+4c)^3(135+8c)(-7+10c)^3(11+14c)$$

$d = 8 : 7$ fields :

$$\mathcal{N}(G, \partial^5 G), \quad \mathcal{N}(\mathcal{N}(L, G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(L, \partial G), \partial^2 G), \\ \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^2 L), \quad \mathcal{N}(L, \partial^4 L) \\ \det D_8^{[1]} = \frac{5184000}{24191167}(-1+c)c^7(11+c)^2(21+4c)^4(135+8c)(-7+10c)^4(11+14c)^2$$

$d = \frac{17}{2} : 4$ fields :

$$\mathcal{N}(L, \partial^5 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 G), \partial G) \\ \det D_{\frac{17}{2}}^{[1]} = \frac{36864}{65065}c^4(11+c)(21+4c)^3(-7+10c)^3(11+14c)$$

$d = 9 : 4$ fields :

$$\mathcal{N}(\mathcal{N}(L, G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(L, \partial G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^3 L) \\ \det D_9^{[1]} = \frac{108}{343}(-1+c)c^4(11+c)(21+4c)^3(-7+10c)^3(11+14c)$$

$d = \frac{19}{2} : 9$ fields :

$$\mathcal{N}(\mathcal{N}(G, \partial G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, G), \partial G), \partial^2 G), \\ \mathcal{N}(L, \partial^6 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^2 G), \\ \mathcal{N}(\mathcal{N}(L, \partial^2 L), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 L), G), \\ \mathcal{N}(\mathcal{N}(L, \partial^4 L), G) \\ \det D_{\frac{19}{2}}^{[1]} = \frac{8599633920}{53202877}(-1+c)^2c^9(11+c)^3(21+4c)^6 \\ (114+5c)(135+8c)(-7+10c)^6(11+14c)^3$$

$d = 10 : 12$ fields :

$$\mathcal{N}(G, \partial^7 G), \quad \mathcal{N}(\mathcal{N}(L, G), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(L, \partial G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 G), \partial^3 G), \\ \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial G), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), G), \partial G), \\ \mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial^2 L), G), \partial G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), L), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^2 L) \\ \mathcal{N}(\mathcal{N}(L, L), \partial^4 L), \quad \mathcal{N}(L, \partial^6 L) \\ \det D_{10}^{[1]} = \frac{1011316948992000000}{762819593107}(-1+c)^2c^{12}(11+c)^4(21+4c)^8(114+5c) \\ (135+8c)^2(-7+10c)^8(11+14c)^4(95+22c)$$

$d = \frac{21}{2} : 10$ fields :

$$\mathcal{N}(\mathcal{N}(G, \partial G), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, G), \partial G), \partial^3 G), \quad \mathcal{N}(L, \partial^7 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^5 G), \\ \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 L), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), \partial G), \\ \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 L), \partial G), \quad \mathcal{N}(\mathcal{N}(L, \partial^4 L), \partial G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^3 L), G) \\ \det D_{\frac{21}{2}}^{[1]} = \frac{477199206497805926400000000}{65518365270271857229}(-1+c)^2c^{10}(11+c)^3(21+4c)^7 \\ (135+8c)(-7+10c)^7(11+14c)^4(95+22c)$$

$d = 11 : 9$ fields :

$$\begin{aligned} & \mathcal{N}(\mathcal{N}(L, G), \partial^6 G), \quad \mathcal{N}(\mathcal{N}(L, \partial G), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 G), \partial^4 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), G), \partial^2 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial^2 L), G), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^3 L), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^5 L) \\ & \det D_{11}^{[1]} = -\frac{28179280429056}{2215547}(-1+c)^2 c^9(11+c)^3(21+4c)^7(135+8c)(-7+10c)^7(11+14c)^4 \end{aligned}$$

$d = \frac{23}{2} : 16$ fields :

$$\begin{aligned} & \mathcal{N}(\mathcal{N}(G, \partial G), \partial^6 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, G), \partial G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, G), \partial^2 G), \partial^3 G) \\ & \mathcal{N}(L, \partial^8 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial G), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^6 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 L), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), \partial^2 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 L), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^4 L), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^3 L), \partial G) \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), L), G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^2 L), G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^4 L), G), \quad \mathcal{N}(\mathcal{N}(L, \partial^6 L), G) \\ & \det D_{\frac{23}{2}}^{[1]} = \frac{16543163447903718821855232000000}{217118368393969}(-1+c)^4 c^{16}(11+c)^6(21+4c)^{12} \\ & (115+4c)(114+5c)(135+8c)^3(-7+10c)^{12}(11+14c)^6(95+22c) \end{aligned}$$

$d = 12 : 23$ fields :

$$\begin{aligned} & \mathcal{N}(\mathcal{N}(\mathcal{N}(G, \partial G), \partial^2 G), \partial^3 G), \quad \mathcal{N}(G, \partial^9 G), \quad \mathcal{N}(\mathcal{N}(L, G), \partial^7 G), \\ & \mathcal{N}(\mathcal{N}(L, \partial G), \partial^6 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 G), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^3 G), \partial^4 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 G), \partial^3 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial G), \partial^2 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial^2 L), G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial^2 L), \partial G), \partial^2 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), G), \partial G), L), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial^2 L), G), \partial G), L), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial^4 L), G), \partial G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), L), L), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), \partial^2 L), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^4 L), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 L), \partial^2 L), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^6 L), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 L), \partial^4 L), \quad \mathcal{N}(L, \partial^8 L) \\ & \det D_{12}^{[1]} = \frac{69893115257789204783084854925269516104499200000000000000000000}{20710629462273412554478537721}(-1+c)^6 c^{23} \\ & (11+c)^8(21+4c)^{16}(115+4c)(114+5c)^2(135+8c)^4(-7+10c)^{17} \\ & (11+14c)^9(95+22c)^2(161+26c)(-81+70c) \end{aligned}$$

$d = \frac{25}{2} : 19$ fields :

$$\begin{aligned} & \mathcal{N}(\mathcal{N}(G, \partial G), \partial^7 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, G), \partial G), \partial^5 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, G), \partial^2 G), \partial^4 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, \partial G), \partial^2 G), \partial^3 G), \\ & \mathcal{N}(L, \partial^9 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), G), \partial G), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(L, L), \partial^7 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^2 L), \partial^5 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), \partial^3 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 L), \partial^3 G), \quad \mathcal{N}(\mathcal{N}(L, \partial^4 L), \partial^3 G), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^3 L), \partial^2 G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), L), L), \partial G), L), \\ & \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^2 L), \partial G), L), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^4 L), \partial G), \\ & \mathcal{N}(\mathcal{N}(L, \partial^6 L), \partial G), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^3 L), G), L), \quad \mathcal{N}(\mathcal{N}(\mathcal{N}(L, L), \partial^5 L), G) \\ & \det D_{\frac{25}{2}}^{[1]} = -\frac{35952348246922073178578114668020015110823318926445772800000000000000000000}{12565355615115842815180391716848789977906099} \\ & (-1+c)^5 c^{19}(11+c)^7(21+4c)^{14}(114+5c)(135+8c)^3(-7+10c)^{15} \\ & (11+14c)^8(95+22c)(161+26c)(-81+70c) \end{aligned}$$

Appendix B: A basis of quasiprimary fields up to dimension $d(\Phi)+6$ built up by one super primary field Φ and $\mathcal{L}$

$d = d(\Phi) : 1 \text{ field} :$

ϕ

$d = d(\Phi) + \frac{1}{2} : 1 \text{ field} :$

ψ

$d = d(\Phi) + \frac{3}{2} : 1 \text{ field} :$

$\mathcal{N}(\phi, G)$

$d = d(\Phi) + 2 : 2 \text{ fields} :$

$\mathcal{N}(\phi, L), \mathcal{N}(\psi, G)$

$d = d(\Phi) + \frac{5}{2} : 2 \text{ fields} :$

$\mathcal{N}(\phi, \partial G), \mathcal{N}(\psi, L)$

$d = d(\Phi) + 3 : 2 \text{ fields} :$

$\mathcal{N}(\phi, \partial L), \mathcal{N}(\psi, \partial G)$

$d = d(\Phi) + \frac{7}{2} : 3 \text{ fields} :$

$\mathcal{N}(\phi, \partial^2 G), \mathcal{N}(\mathcal{N}(\phi, L), G), \mathcal{N}(\psi, \partial L)$

$d = d(\Phi) + 4 : 5 \text{ fields} :$

$\mathcal{N}(\mathcal{N}(\phi, G), \partial G), \mathcal{N}(\mathcal{N}(\phi, L), L), \mathcal{N}(\phi, \partial^2 L),$
 $\mathcal{N}(\psi, \partial^2 G), \mathcal{N}(\mathcal{N}(\psi, L), G)$

$d = d(\Phi) + \frac{9}{2} : 6 \text{ fields} :$

$\mathcal{N}(\phi, \partial^3 G), \mathcal{N}(\mathcal{N}(\phi, L), \partial G), \mathcal{N}(\mathcal{N}(\phi, \partial L), G),$
 $\mathcal{N}(\mathcal{N}(\psi, G), \partial G), \mathcal{N}(\mathcal{N}(\psi, L), L), \mathcal{N}(\psi, \partial^2 L)$

$d = d(\Phi) + 5 : 6 \text{ fields} :$

$\mathcal{N}(\mathcal{N}(\phi, G), \partial^2 G), \mathcal{N}(\mathcal{N}(\phi, L), \partial L), \mathcal{N}(\phi, \partial^3 L),$
 $\mathcal{N}(\psi, \partial^3 G), \mathcal{N}(\mathcal{N}(\psi, L), \partial G), \mathcal{N}(\mathcal{N}(\psi, \partial L), G)$

$d = d(\Phi) + \frac{11}{2} : 8 \text{ fields} :$

$\mathcal{N}(\phi, \partial^4 G), \mathcal{N}(\mathcal{N}(\phi, L), \partial^2 G), \mathcal{N}(\mathcal{N}(\phi, \partial L), \partial G),$
 $\mathcal{N}(\mathcal{N}(\mathcal{N}(\phi, L), L), G), \mathcal{N}(\mathcal{N}(\phi, \partial^2 L), G), \mathcal{N}(\mathcal{N}(\psi, G), \partial^2 G),$
 $\mathcal{N}(\mathcal{N}(\psi, L), \partial L), \mathcal{N}(\psi, \partial^3 L)$

$d = d(\Phi) + 6 : 12 \text{ fields} :$

$\mathcal{N}(\mathcal{N}(\phi, G), \partial^3 G), \mathcal{N}(\mathcal{N}(\phi, \partial G), \partial^2 G), \mathcal{N}(\mathcal{N}(\mathcal{N}(\phi, L), G), \partial G),$
 $\mathcal{N}(\mathcal{N}(\mathcal{N}(\phi, L), L), L), \mathcal{N}(\mathcal{N}(\phi, L), \partial^2 L), \mathcal{N}(\mathcal{N}(\phi, \partial L), \partial L),$
 $\mathcal{N}(\phi, \partial^4 L), \mathcal{N}(\psi, \partial^4 G), \mathcal{N}(\mathcal{N}(\psi, L), \partial^2 G),$
 $\mathcal{N}(\mathcal{N}(\psi, \partial L), \partial G), \mathcal{N}(\mathcal{N}(\mathcal{N}(\psi, L), L), G), \mathcal{N}(\mathcal{N}(\psi, \partial^2 L), G)$

**Appendix C: Determinants of quasiprimary parts of the Kac-matrix
with one additional super primary field involved**

$SW(\frac{3}{2}, 2) :$

$$\det D_{\frac{7}{2}}^{[2]} = \frac{1}{15}c(6 + 5c)$$

$$\det D_4^{[2]} = -\frac{1}{150}c^2(29 + 2c)(6 + 5c)$$

$SW(\frac{3}{2}, \frac{7}{2}) :$

$$\det D_5^{[\frac{7}{2}]} = -\frac{1}{21}c(21 + 4c)$$

$$\det D_{\frac{11}{2}}^{[\frac{7}{2}]} = \frac{1}{336}c^2(53 + 2c)(21 + 4c)$$

$$\det D_6^{[\frac{7}{2}]} = \frac{1}{40}(-1 + c)c^2(53 + 2c)$$

$$\det D_{\frac{13}{2}}^{[\frac{7}{2}]} = \frac{4}{121}(-1 + c)c^2(21 + 4c)$$

$SW(\frac{3}{2}, 4) :$

$$\det D_{\frac{11}{2}}^{[4]} = \frac{1}{18}c(20 + 3c)$$

$$\det D_6^{[4]} = -\frac{1}{324}c^2(61 + 2c)(20 + 3c)$$

$$\det D_{\frac{13}{2}}^{[4]} = -\frac{1}{495}c^2(61 + 2c)(-7 + 10c)$$

$$\det D_7^{[4]} = -\frac{1}{270}c^2(20 + 3c)(-7 + 10c)$$

$$\det D_{\frac{15}{2}}^{[4]} = -\frac{1}{16731}c^3(61 + 2c)(20 + 3c)^2(-65 + 44c)$$

$$\det D_8^{[4]} = \frac{1}{29513484}c^5(5 + 2c)(61 + 2c)^2(20 + 3c)^2(-7 + 10c)(377 + 10c)(-65 + 44c)$$

$SW(\frac{3}{2}, \frac{11}{2}) :$

$$\det D_7^{[\frac{11}{2}]} = -\frac{4}{33}c(11 + c)$$

$$\det D_{\frac{15}{2}}^{[\frac{11}{2}]} = \frac{1}{198}c^2(11 + c)(85 + 2c)$$

$$\det D_8^{[\frac{11}{2}]} = \frac{1}{1092}c^2(85 + 2c)(5 + 13c)$$

$$\det D_{\frac{17}{2}}^{[\frac{11}{2}]} = \frac{16}{2925}c^2(11 + c)(5 + 13c)$$

$$\det D_9^{[\frac{11}{2}]} = -\frac{2}{1155}c^3(11 + c)^2(85 + 2c)(-10 + 7c)$$

$$\det D_{\frac{19}{2}}^{[\frac{11}{2}]} = \frac{1}{2203047}c^5(11 + c)^2(85 + 2c)^2(-10 + 7c)(5 + 13c)(4717 + 1092c + 20c^2)$$

$$\det D_{10}^{[\frac{11}{2}]} = \frac{5}{20857419}c^6(11 + c)^2(15 + c)(85 + 2c)^2(-5 + 4c)(5 + 13c)^2(4717 + 1092c + 20c^2)$$

$$\det D_{\frac{21}{2}}^{[\frac{11}{2}]} = \frac{819200}{19321000497}c^6(11 + c)^3(15 + c)(85 + 2c)^2(-5 + 4c)(-10 + 7c)(5 + 13c)^2$$

$SW(\frac{3}{2}, 6) :$

$$\det D_{\frac{15}{2}}^{[6]} = \frac{1}{117}c(162 + 13c)$$

$$\det D_8^{[6]} = -\frac{1}{3042}c^2(93 + 2c)(162 + 13c)$$

$$\det D_{\frac{17}{2}}^{[6]} = -\frac{1}{1365}c^2(93 + 2c)(11 + 14c)$$

$$\det D_9^{[6]} = -\frac{1}{2912}c^2(162 + 13c)(11 + 14c)$$

$$\begin{aligned}
det D_{\frac{19}{2}}^{[6]} &= -\frac{1}{338130}c^3(93+2c)(162+13c)^2(-27+20c) \\
det D_{10}^{[6]} &= \frac{1}{1602229005}c^5(93+2c)^2(162+13c)^2(11+14c)(-27+20c)(5917+1188c+20c^2) \\
det D_{\frac{21}{2}}^{[6]} &= -\frac{16}{186752880639}c^6(33+2c)(93+2c)^2(162+13c)^2(11+14c)^2 \\
&\quad (-27+20c)(5917+1188c+20c^2) \\
det D_{11}^{[6]} &= -\frac{2}{4461283125}c^6(33+2c)(93+2c)^2(162+13c)^3(11+14c)^2(-27+20c)^2 \\
det D_{\frac{23}{2}}^{[6]} &= -\frac{2048}{86728774951175175}c^8(93+2c)^3(162+13c)^5(11+14c)^2 \\
&\quad (-27+20c)^2(-43+85c)(5917+1188c+20c^2) \\
det D_{12}^{[6]} &= -\frac{8192}{412159655278351185230025}c^{12}(33+2c)(93+2c)^5(162+13c)^6(11+14c)^3 \\
&\quad (-27+20c)^3(-43+85c)(5917+1188c+20c^2)^2(-837+8540c+140c^2)
\end{aligned}$$

Appendix D: Realization of $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ at $c = \frac{10}{7}$ as a subalgebra of $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$

As mentioned in chapter four and five there is a strange connection between these algebras at $c = \frac{10}{7}$. $\mathcal{SW}(\frac{3}{2}, \frac{5}{2})$ contains a super primary field of dimension $\frac{11}{2}$. The two components of this field are:

$$\begin{aligned}
\tilde{\phi} &= \alpha \left(\mathcal{N}(\psi, \phi) + \frac{8869}{29580}C_{G\phi}^{\psi}\mathcal{N}(L, \partial^2 G) - \frac{5537}{25143}C_{G\phi}^{\psi}\mathcal{N}(\mathcal{N}(L, L), G) \right) \\
\tilde{\psi} &= \frac{\alpha}{\sqrt{2}} \left(\mathcal{N}(\psi, \psi) + \frac{1}{6}\mathcal{N}(\phi, \partial\phi) + \frac{8575}{67048}\mathcal{N}(G, \partial^3 G) + \right. \\
&\quad \left. \frac{206143}{251430}\mathcal{N}(L, \partial^2 L) - \frac{5537}{25143}\mathcal{N}(\mathcal{N}(L, G), \partial G) - \frac{11074}{25143}\mathcal{N}(\mathcal{N}(L, L), L) \right) \\
\alpha^2 &= \frac{176001}{1567918}
\end{aligned}$$

where $\Phi = \phi + \theta\psi$ is the simple super primary field of dimension $\frac{5}{2}$. The fields $\tilde{\phi}, \tilde{\psi}$ are orthogonal to all other fields of dimension $\frac{11}{2}, 6$ and their normalization is described by (3.3'). We have calculated the self-coupling:

$$(C_{\tilde{\phi}\tilde{\phi}}^{\tilde{\psi}})^2 = \frac{844918800}{34554863}$$

This is exactly the coupling obtained in the direct construction of $\mathcal{SW}(\frac{3}{2}, \frac{11}{2})$ at $c = \frac{10}{7}$. The two different signs of the coupling are realized by the two possible signs for α .

References

- [1] A.A. Belavin, A.M. Polyakov, A.B. Zamolodchikov, Nucl. Phys. **B241** (1984) p. 333
- [2] A.B. Zamolodchikov, Theor. Math. Phys. **65** (1986) p. 1205
- [3] F.A. Bais, P. Bouwknegt, M. Surridge, K. Schoutens, Nucl. Phys. **B304** (1988) p. 371
- [4] A. Bilal, J.L. Gervais, Nucl. Phys. **B318** (1989) p. 579
- [5] J. Balog, L. Fehér, P. Forgács, L. O’Raifeartaigh, A. Wipf
Phys. Lett. **B244** (1990) p. 435
- [6] K. Hamada, M. Takao, Phys. Lett. **B209** (1988) p. 247
Erratum, Phys. Lett. **B213** (1988) p. 564
- [7] D.H. Zhang, Phys. Lett. **B232** (1989) p. 323
- [8] J.M. Figueroa-O’Farrill, S. Schrans, Phys. Lett. **B245** (1990) p. 471
- [9] P. Bouwknegt, Phys. Lett. **B207** (1988) p. 295
- [10] R. Blumenhagen, M. Flohr, A. Kliem, W. Nahm, A. Recknagel, R. Varnhagen
Nucl. Phys. **B361** (1991) p. 255
- [11] H.G. Kausch, G.M.T. Watts, Nucl. Phys. **B354** (1991) p. 740
- [12] W. Nahm, in proceedings Trieste Conference on
‘Recent Developments in Conformal Field Theories’, ICTP, Trieste (1989) p. 81
- [13] D. Friedan, Z. Qiu, S. Shenker, Phys. Lett. **B151** (1985) p. 37
- [14] T. Inami, Y. Matsuo, I. Yamanaka, Phys. Lett. **B215** (1988) p. 701
- [15] S. Komata, K. Mohri, H. Nohara, Nucl. Phys. **B359** (1991) p. 168
- [16] K. Hornfeck, E. Ragoucy, Nucl. Phys. **B340** (1990) p. 225
- [17] J.M. Figueroa-O’Farrill, S. Schrans, Int. Jour. of Mod. Phys. **A7** (1992) p. 591
- [18] R. Blumenhagen, preprint BONN-HE-91-20 to be published in Nucl. Phys. **B**
- [19] K. Hornfeck, preprint London, King’s College 91.08.26
- [20] W. Nahm, *Conformal Quantum Field Theories in Two Dimensions*
World Scientific, to be published
- [21] P. West, *Introduction to Supersymmetry and Supergravity*
World Scientific (1986)
- [22] M. Flohr, Diplomarbeit BONN-IR-91-30
- [23] P. Bouwknegt, in proceedings of the meeting ‘Infinite dimensional Lie algebras and Groups’
C.I.R.M., Luminy, Marseille (1988) p. 527
- [24] A. Cappelli, C. Itzykson, J.B. Zuber, Comm. Math. Phys. **113** (1987) p. 1
- [25] H.G. Kausch, Phys. Lett. **B259** (1991) p. 448
- [26] M. Flohr, preprint BONN-HE-92-08
- [27] W. Eholzer, M. Flohr, A. Honecker, R. Hübel, W. Nahm, R. Varnhagen
preprint BONN-HE-91-22 to be published in Nucl. Phys. **B**
- [28] M.A. Bershadsky, V.G. Knizhnik, M.G. Teitelman, Phys. Lett. **B151** (1985) p. 31
- [29] H. Eichenherr, Phys. Lett. **B151** (1985) p. 26
- [30] G. Mussardo, G. Sotkov, M. Stanishkov, Phys. Lett. **B195** (1987) p. 397,
Nucl. Phys. **B305** (1988) p.69
- [31] P. Goddard, A. Kent, D. Olive, Comm. Math. Phys. **103** (1986) p. 105
- [32] A.N. Schellekens, S. Yankielowicz, Nucl. Phys. **B327** (1989) p. 673
- [33] S. Doplicher, R. Haag, J.E. Roberts, Comm. Math. Phys. **23** (1971) p. 199,
Comm. Math. Phys. **35** (1974) p. 49

- [34] K. Fredenhagen, K.H. Rehren, B. Schroer, Comm. Math. Phys. 125 (1989) p. 201
- [35] F. Wilczek, Phys. Rev. Lett. 49 (1982) p. 957
- [36] W. Eholzer, A. Honecker, R. Hübel
Representations of $N = 1$ Extended Superconformal Algebras , in preparation
- [37] F. Delduc, E. Ragoucy, P. Sorba, preprint Lyon, ENSLAPP-L-352-91
- [38] K. Schoutens, A. Sevrin, Phys. Lett. **B258** (1991) p. 134
- [39] R. Varnhagen, Phys. Lett. **B275** (1992) p. 87